

Flow equivalence of shift spaces (and their C^* -algebras), I

Søren Eilers

eilers@math.ku.dk

Department of Mathematical Sciences
University of Copenhagen

18.01.11


Content

- 1 Definitions
- 2 Conjugacy
- 3 Classification
- 4 Flow equivalence
- 5 Flow classification

Outline


- 1 Definitions
- 2 Conjugacy
- 3 Classification
- 4 Flow equivalence
- 5 Flow classification

Baker's map



101110100101001111100...

Irrational rotation



0001000100100010010001000...

Symbolic dynamics

Let \mathfrak{a} be a finite set and equip $\mathfrak{a}^{\mathbb{Z}}$ with the product topology based on the discrete topology on \mathfrak{a} .

Definition

A **shift space** is a subset X of $\mathfrak{a}^{\mathbb{Z}}$ which is closed and closed under the **shift map**

$$\sigma : \mathfrak{a}^{\mathbb{Z}} \rightarrow \mathfrak{a}^{\mathbb{Z}} \quad \sigma((x_i)) = (x_{i+1})$$

Definition

A shift space is **irreducible** if some forward orbit $\{\sigma^n(x) \mid n \in \mathbb{N}\}$ is dense.

3 constructions

Name	Input	Description	Example
$X^{(W)}$	List of words W	Sequences not containing words from W	$W = \{11\}$
X_G	Graph G	Infinite paths on G	
$L_{\mathcal{A}}$	Labelled graph \mathcal{A}	Infinite paths on \mathcal{A}	

Forbidden word shifts

Let W be a set of finite words on \mathfrak{a} .

Definition

$X^{(W)}$ is the shift space $\{x \in \mathfrak{a}^{\mathbb{Z}} \mid \forall i < j : x_i \cdots x_j \notin W\}$

Example

With $\mathfrak{a} = \{0, 1\}$ and $W = \{11\}$ the shift space $X^{(W)}$ contains elements such as

$\cdots 01000010001000100001010101001001000100010 \cdots$

Lemma

For any shift space X , $X = X^{(W)}$ where W is chosen as the complement of the language

$$\mathcal{L}(X) = \{x_i \cdots x_j \mid x \in X, i < j\}$$

Edge shifts

Let a graph $G = (V, E, r, s)$ be given with


- Vertices V
- Edges E enumerated $\{e_1, \dots, e_n\}$
- Range and source maps $r, s : E \rightarrow V$.

Definition

X_G is the shift space $X^{(W)}$ with alphabet E and

$$W = \{e_i e_j \mid r(e_i) \neq s(e_j)\}$$

Example

With $G =$  , X_G contains elements such as

$\dots e_1 e_1 e_2 e_3 e_2 e_3 e_2 e_3 e_1 e_2 e_3 e_2 e_3 e_1 e_2 e_3 e_2 e_3 e_1 e_1 e_1 e_1 e_1 e_2 \dots$

Labelled edge shifts

Convention

A **labelled graph** $\mathcal{A} = (V, E, r, s, \mathfrak{a}, \lambda)$ is given by an underlying graph (V, E, r, s) and a labelling map $\Lambda : E \rightarrow \mathfrak{a}$

Definition

We denote by $X_{\mathcal{A}}$ the edge shift associated to the underlying graph of \mathcal{A} and by

$$\lambda : X_{\mathcal{A}} \rightarrow \mathfrak{a}^{\mathbb{Z}}$$

the *labelling map* induced by Λ . The shift defined by \mathcal{A} is $L_{\mathcal{A}} = \lambda(X_{\mathcal{A}})$.

Labelled edge shifts

Example

With $\mathcal{A} = 0 \begin{array}{c} \curvearrowright \\ \bullet \end{array} \begin{array}{c} \xrightarrow{1} \\ \xleftarrow{0} \\ \bullet \end{array}$ the shift space $X_{\mathcal{A}}$ contains elements
such as

$\dots 01000010001000100001010101001001000100010 \dots$

Outline

- 1 Definitions
- 2 Conjugacy**
- 3 Classification
- 4 Flow equivalence
- 5 Flow classification

Definition

Let $X \subseteq \mathfrak{a}^{\mathbb{Z}}$ and $Y \subseteq \mathfrak{b}^{\mathbb{Z}}$. $\phi : X \rightarrow Y$ is the (m, n) *sliding block code* given by a map

$$\Phi : \mathfrak{a}^{n+1+m} \rightarrow \mathfrak{b}$$

when

$$\phi(x)_i = \Phi(x_{i-m} \cdots x_{i+n})$$

Lemma

The following are equivalent:

- ϕ is continuous and shift-commuting
- ϕ is a sliding block code

Definition

X and Y are *conjugate* when there is a bijective sliding block code $\phi : X \rightarrow Y$

Shifts of finite type

Definition

A shift space is a *shift of finite type (SFT)* if it has the form $X^{(W)}$ with W finite.

Lemma

The following are equivalent:

- X is an SFT
- $X \simeq X_G$ for some graph G

Sofic shifts

Definition

A shift space is *sofic* if there is a surjective sliding block code $\phi : Y \rightarrow X$ with Y an SFT.

Lemma

The following are equivalent:

- X is sofic
- $X \simeq L_{\mathcal{A}}$ for some labelled graph \mathcal{A}

Theorem

When X is irreducible and sofic, there is a unique labelled graph \mathcal{A} with fewest possible vertices and each pair of edges emanating from the same vertex distinctly labelled, such that $X \simeq L_{\mathcal{A}}$. \mathcal{A} is called the Fischer cover of X .

Outline

- 1 Definitions
- 2 Conjugacy
- 3 Classification**
- 4 Flow equivalence
- 5 Flow classification

Classification



The classification problem

Let X and Y be shift spaces finitely presented by objects \mathcal{A} and \mathcal{B} , respectively. Determine in terms of \mathcal{A} and \mathcal{B} when X and Y are conjugate.

The SFT classification problem

Let X and Y be irreducible shifts of finite type given by graphs G and H , respectively. Determine in terms of G and H when X and Y are conjugate.

State splitting



$$\begin{bmatrix} 2 & 1 \\ 2 & 0 \end{bmatrix} = \begin{bmatrix} 2 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 2 & 0 & 1 \\ 2 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 2 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix}$$

Theorem (Williams)

Let X_G and X_H be two irreducible SFTs given by graphs with adjacency matrices A and B , respectively. The following conditions are equivalent.





- (i) X_G and X_H are conjugate.
- (ii) There exist nonnegative integral matrices D_i and E_i with

$$A = D_0 E_0, E_0 D_0 = D_1 E_1, \dots, E_n D_n = B$$





Arsenal of invariants

Real numbers (entropy), power series (zeta function), ordered abelian groups (Dimension group), finitely generated abelian groups (Bowen-Franks groups), C^* -algebras (Cuntz-Krieger algebra),...

4 examples

A	G	$h(X_G)$	$BF(X_G)$
$\begin{bmatrix} 2 & 2 \\ 2 & 2 \end{bmatrix}$		4	$(\mathbb{Z}_3, -)$
$\begin{bmatrix} 3 & 1 \\ 3 & 1 \end{bmatrix}$		4	$(\mathbb{Z}_3, -)$
$\begin{bmatrix} 1 & 1 \\ 3 & 2 \end{bmatrix}$		$\frac{3+\sqrt{13}}{2}$	$(\mathbb{Z}_3, -)$
$\begin{bmatrix} 2 & 2 \\ 1 & 3 \end{bmatrix}$		4	$(\mathbb{Z}, 0)$

4 examples

A	G	$h(X_G)$	$BF(X_G)$
$\begin{bmatrix} 2 & 2 \\ 2 & 2 \end{bmatrix}$		4	$(\mathbb{Z}_3, -)$
$\begin{bmatrix} 3 & 1 \\ 3 & 1 \end{bmatrix}$		4	$(\mathbb{Z}_3, -)$
$\begin{bmatrix} 1 & 1 \\ 3 & 2 \end{bmatrix}$		$\frac{3+\sqrt{13}}{2}$	$(\mathbb{Z}_3, -)$
$\begin{bmatrix} 2 & 2 \\ 1 & 3 \end{bmatrix}$		4	$(\mathbb{Z}, 0)$

Outline

- 1 Definitions
- 2 Conjugacy
- 3 Classification
- 4 Flow equivalence**
- 5 Flow classification

Flow equivalence

Associated to any shift space there is a **suspension flow** given by product topology on

$$SX = \frac{X \times \mathbb{R}}{(x, t) \sim (\sigma(x), t + 1)}$$

Definition

X and Y are *flow equivalent* (written $X \simeq_{fe} Y$) when SX and SY are homeomorphic in a way preserving direction in \mathbb{R} .

Theorem (Parry-Sullivan)

Flow equivalence is the coarsest equivalence relation containing conjugacy and $X \sim X_{a \rightarrow a^}$*

Flow classification

Lemma

If $X \simeq_{fe} Y$ and X is SFT, sofic or irreducible, then so is Y .

The flow classification problem

Let X and Y be shifts finitely presented by objects \mathcal{A} and \mathcal{B} , respectively. Determine in terms of \mathcal{A} and \mathcal{B} when X and Y are flow equivalent.

The SFT flow classification problem

Let X and Y be irreducible shifts of finite type given by graphs G and H , respectively. Determine in terms of G and H when X and Y are flow equivalent.

Outline

- 1 Definitions
- 2 Conjugacy
- 3 Classification
- 4 Flow equivalence
- 5 Flow classification**

Flow classification of SFTs

Theorem (Franks)

Let X_G and X_H be two irreducible SFTs given by graphs with adjacency matrices A and B , respectively. The following conditions are equivalent.

(i) $X_G \simeq_{fe} X_H$





(ii)

$$\mathbb{Z}^m / (1 - A)\mathbb{Z}^m \simeq \mathbb{Z}^n / (1 - B)\mathbb{Z}^n$$





and

$$\operatorname{sgn} \det(1 - A) = \operatorname{sgn} \det(1 - B)$$

4 examples

A	G	$BF(X_G)$
$\begin{bmatrix} 2 & 2 \\ 2 & 2 \end{bmatrix}$		$(\mathbb{Z}_3, -)$
$\begin{bmatrix} 3 & 1 \\ 3 & 1 \end{bmatrix}$		$(\mathbb{Z}_3, -)$
$\begin{bmatrix} 1 & 1 \\ 3 & 2 \end{bmatrix}$		$(\mathbb{Z}_3, -)$
$\begin{bmatrix} 2 & 2 \\ 1 & 3 \end{bmatrix}$		$(\mathbb{Z}, 0)$

4 examples

A	G	$BF(X_G)$
$\begin{bmatrix} 2 & 2 \\ 2 & 2 \end{bmatrix}$		$(\mathbb{Z}_3, -)$
$\begin{bmatrix} 3 & 1 \\ 3 & 1 \end{bmatrix}$		$(\mathbb{Z}_3, -)$
$\begin{bmatrix} 1 & 1 \\ 3 & 2 \end{bmatrix}$		$(\mathbb{Z}_3, -)$
$\begin{bmatrix} 2 & 2 \\ 1 & 3 \end{bmatrix}$		$(\mathbb{Z}, 0)$

Flow classification of sofic

Theorem

Let X and Y be two irreducible sofic shifts and let \mathcal{A}, \mathcal{B} be their Fischer covers. The following conditions are equivalent.

(i) $X \simeq_{fe} Y$

(ii)
$$\begin{array}{ccc} SX_{\mathcal{A}} & \xrightarrow{\sim+} & SX_{\mathcal{B}} \\ S\lambda_{\mathcal{A}} \downarrow & & \downarrow S\lambda_{\mathcal{B}} \\ SL_{\mathcal{A}} & \xrightarrow{\sim+} & SL_{\mathcal{B}} \end{array}$$

Multiplicity set

Definition

With a given map $\lambda : X_{\mathcal{A}} \rightarrow L_{\mathcal{A}}$ we set

$$\begin{aligned}\widetilde{L}_{\mathcal{A}} &= \{x \in L_{\mathcal{A}} \mid |\lambda^{-1}(\{x\})| > 1\} \\ \widetilde{X}_{\mathcal{A}} &= \lambda^{-1}(\widetilde{L}_{\mathcal{A}})\end{aligned}$$

and restrict λ to

$$\widetilde{\lambda} : \widetilde{X}_{\mathcal{A}} \rightarrow \widetilde{L}_{\mathcal{A}}$$

Example

With $\mathcal{A} = 0 \begin{array}{c} \circlearrowleft \\ \bullet \end{array} \begin{array}{c} \xrightarrow{1} \\ \bullet \\ \xleftarrow{0} \end{array}$ and $\mathcal{B} = 1 \begin{array}{c} \circlearrowleft \\ \bullet \end{array} \begin{array}{c} \xrightarrow{0} \\ \bullet \\ \xleftarrow{0} \end{array}$ we get $\widetilde{L}_{\mathcal{A}} = \emptyset$
and $\widetilde{L}_{\mathcal{B}} = \{0^\infty\}$.

Theorem (Boyle-Carlsen-E)

Let X and Y be two irreducible sofic shift spaces with Fischer covers \mathcal{A} and \mathcal{B} , respectively, and assume that $\widetilde{X}_{\mathcal{A}}$ and $\widetilde{X}_{\mathcal{B}}$ are both closed. Then X and Y are flow equivalent exactly when the following conditions hold:

$$(1) \quad X_{\mathcal{A}} \simeq_{fe} X_{\mathcal{B}}$$

$$(2) \quad \begin{array}{ccc} S\widetilde{X}_{\mathcal{A}} & \xrightarrow{\sim_+} & S\widetilde{X}_{\mathcal{B}} \\ S\widetilde{\lambda}_{\mathcal{A}} \downarrow & & \downarrow S\widetilde{\lambda}_{\mathcal{B}} \\ S\widetilde{L}_{\mathcal{A}} & \xrightarrow{\sim_+} & S\widetilde{L}_{\mathcal{B}} \end{array}$$

$\lambda : X_{\mathcal{A}} \rightarrow L_{\mathcal{A}}$	$\tilde{\lambda} : \tilde{X}_{\mathcal{A}} \rightarrow \tilde{L}_{\mathcal{A}}$

$\lambda : X_{\mathcal{A}} \rightarrow L_{\mathcal{A}}$	$\tilde{\lambda} : \tilde{X}_{\mathcal{A}} \rightarrow \tilde{L}_{\mathcal{A}}$