

Weights of partitions and character zeros

Christine Bessenrodt, Jørn B. Olsson*

June 23, 2004

Abstract

Motivated by a question on the zeros of irreducible characters of the symmetric and alternating groups, we classify partitions which are of maximal p -weight for all odd primes p .

1 Introduction

A well-known result by Burnside states that any non-linear character of a finite group vanishes on some element of the group. Recently this was refined in [5], where it was shown that such a character always has a zero at an element of prime power order. Moreover, it had been noticed in [5] that any non-linear character of a finite simple group except possibly the alternating groups even vanishes on some element of prime order. Here we show that this character property also holds for the alternating and the symmetric groups. Indeed, this vanishing property is a simple consequence of a combinatorial result on the weights of partitions, which may be of independent interest.

The elements of prime order p which we are going to use in the symmetric group S_n are those of maximal weight, i.e., they are a product of $\lfloor \frac{n}{p} \rfloor$ p -cycles. (Here $\lfloor \cdot \rfloor$ denotes the floor function. Thus $\lfloor x \rfloor$ is the integral part of $x \in \mathbb{R}$.)

Consider a partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_m)$ of the integer n . For a given integer $r \in \mathbb{N}$, we denote by $w_r(\lambda)$ the r -weight of λ , i.e., $w_r(\lambda)$ is the maximal number of r -hooks that can successively be removed from λ . The resulting partition after removing this maximal number of r -hooks is then

⁰MSC Classification: 20C30, 05A17

*Partially supported by The Danish National Research Council.

the r -core $\lambda_{(r)}$ of λ (see [4] or [6] for details). In this paper, we will only deal with the case where r is a prime number p . We present a classification of the partitions which have maximal p -weight $\left\lfloor \frac{n}{p} \right\rfloor$ for all (odd) primes p .

The relevance of this classification to the question about the vanishing of character values is easily explained: The irreducible characters of S_n are labelled by the partitions λ of n . The Murnaghan-Nakayama formula for character values in S_n shows that the irreducible character labelled by λ vanishes on a p -element of maximal weight, if the p -weight of λ is *not* maximal.

Our main result is then the following:

Theorem 1.1 *Let λ be a partition of $n \in \mathbb{N}$. Then the following holds:*

(1) *λ is of maximal p -weight for all primes p , if and only if one of the following occurs:*

$$\lambda = (n), (1^n) \text{ or } (2^2).$$

(2) *λ is of maximal p -weight for all primes $p > 2$, if and only if λ is one of the partitions in (1), one of $(n-1, 1)$, $(2, 1^{n-2})$, where $n = 2^a + 1$ for some $a \in \mathbb{N}$, or one of the following occurs:*

$$\begin{aligned} n = 6 : \lambda &= (3, 2, 1) \\ n = 8 : \lambda &= (5, 2, 1) \text{ or } (3, 2, 1^3) \\ n = 9 : \lambda &= (6, 3) \text{ or } (2^3, 1^3) \\ n = 10 : \lambda &= (4, 3, 2, 1) \end{aligned}$$

This combinatorial classification result has the following consequence:

Theorem 1.2 *Let $n \in \mathbb{N}$. Let χ be any non-linear character of the symmetric group S_n or the alternating group A_n . Then χ vanishes on some element of prime order. If $\chi(1)$ is not a 2-power, then χ is zero on some element of odd prime order.*

Remark 1.3 If an irreducible character χ of a finite group G has a zero at an element of prime order p , then p divides $\chi(1)$; thus we can only expect zeros of odd prime order if $\chi(1)$ is not a 2-power. Note that the irreducible characters of S_n and A_n of prime power degree have been classified in [1]; from Theorem 1.1 we can immediately recover the classification of irreducible characters of 2-power degree for these groups.

The converse of the statement above does not hold, even for $G = S_n$. We denote by $[\lambda]$ the irreducible character of S_n associated to the partition λ .

Then $[5, 2](1) = 14$, but $[5, 2]$ does not vanish on any element of order 2. This is not just an accident for the prime 2: $[8, 1^3](1) = 120$, but $[8, 1^3]$ has no zero on elements of order 2 or 3.

Theorem 1.2 then allows us to settle the question left open in [5]. Combined with corresponding results on sporadic groups [5, Theorem 3.4] and simple groups of Lie type [5, Theorem 5.1] we obtain:

Corollary 1.4 *Any non-trivial irreducible character of a finite simple group vanishes on some element of prime order.*

We refer to [4, section 2.5] for the labelling of the irreducible characters of A_n . Now [4, 6.2.45] gives a simple relation between the p -weight of a partition λ and the defect of the p -block containing the irreducible character labelled by λ . Therefore the following is another consequence of Theorem 1.1:

Theorem 1.5 (1) *The characters $[n], [1^n]$ and $[2^2]$ are the only irreducible characters of S_n which are in p -blocks of maximal defect for all primes p . Apart from $[1^2], [1^3], [1^4], [1^6], [2^2]$, the trivial character of S_n is the only irreducible character which is in the principal p -block for all primes $p \leq n$.*
 (2) *The characters $\{n\}, \{2, 1\}_\pm$ and $\{2^2\}_\pm$ are the only irreducible characters of A_n which are in p -blocks of maximal defect for all primes p . They belong to the principal p -block for all primes $p \leq n$, except for the characters $\{2, 1\}_\pm$ at $p = 2$.*

The paper is organized as follows. Section 2 contains some results on hook lengths and on large prime divisors of products of consecutive integers. In section 3 we consider partitions of maximal weight and modify an algorithm of [1] to suit our purposes better. The algorithms are used to generate “large” first column hook lengths in the partitions under consideration. The final section contains the proofs of all the results stated above.

2 Preliminaries

We refer to [4], [6] for details about partitions, Young diagrams and hooks. Consider a partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_l)$ of the integer n . Thus $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_l > 0$ and $\lambda_1 + \lambda_2 + \dots + \lambda_l = n$. We call the λ_i 's the parts of λ and $l = l(\lambda)$ the length of λ . Moreover for $i \geq 1$, $m_i = m_i(\lambda)$ denotes the number of parts equal to i in λ . The Young diagram of λ consists of n nodes (boxes) with λ_i nodes in the i th row. We refer to the nodes in matrix notation, i.e.

the (i, j) -node is the j th node in the i th row. The (i, j) -hook consists of the nodes in the Young diagram to the right and below the (i, j) -node, and including this node. The number of nodes in this hook is its hooklength, denoted by h_{ij} . For the hooklengths in the first column we write $h_i = h_{i1}$, $i = 1, \dots, l$. We call any such number h_i an **fch** (first column hooklength) of λ .

We will often make use of the following property of the weight of a partition (see [4, 2.7.37], [6, (3.3)]).

Lemma 2.1 *If λ is a partition of p -weight $w_p(\lambda) = w$, then λ has exactly w hooks of length divisible by p . In particular, if λ has a hook of length divisible by p , then it has a hook of length p .*

We also recall some elementary results about hook lengths from [1, section 2].

Proposition 2.2 *Let λ be a partition of n , $l = l(\lambda)$. Let $1 \leq i < j \leq l$. Then*

$$h_i + h_j \leq n + 1 + m_1 .$$

Lemma 2.3 *Let λ be a partition of n . Suppose that $s = h_{ik}$ and $t = h_{jm}$ where $(i, k) \neq (j, m)$.*

- (1) *If $i \neq j$ and $k \neq m$, then $s + t \leq n$.*
- (2) *If $s + t > n$, then either $i = j = 1$ (both hooks in the first row) or $k = m = 1$ (both hooks in the first column).*

Corollary 2.4 *Let λ be a partition of n . For $j \geq 2$, every hook of length $t > n - h_j = n - h_{j1}$ is a **fch** of λ .*

We will also need some number-theoretic results. Improving on a result by Sylvester and Schur, Hanson [2] proved the following:

Theorem 2.5 *The product of k consecutive numbers all greater than k contains a prime divisor greater than $\frac{3}{2}k$, with the only exceptions $3 \cdot 4$, $8 \cdot 9$ and $6 \cdot 7 \cdot 8 \cdot 9 \cdot 10$.*

Lemma 2.6 *Let $1 < k \leq 500$, $k \neq 3$ or 4 . Then any product of k consecutive integers larger than 825.000 has a prime divisor $q > 2k$.*

Proof. For $k = 2$ the result already follows from Theorem 2.5. So we may assume that $k \geq 5$.

Let $n \in \mathbb{N}$ with $n > 825000$. Assume that all prime factors in the k consecutive numbers $n - k + 1, \dots, n - 1, n$ are at most $2k$.

With $\pi(x) = \#\{p \text{ prime} \mid p \leq x\}$, for $x \in \mathbb{R}_+$, we obtain

$$\left(\frac{n}{k}\right)^k < \binom{n}{k} = \prod_{p \mid \binom{n}{k}} p^{b_p} \leq n^{\pi(2k)}$$

where p^{b_p} is the maximal power of p dividing $\binom{n}{k}$ (this is known to be bounded by n). Equivalently,

$$(k - \pi(2k)) \ln(n) < k \ln k.$$

Note that for $5 \leq k \leq 500$, $k > \pi(2k)$ (this may be checked directly). Hence, if the assumption holds then

$$\ln(n) < \frac{k \ln(k)}{k - \pi(2k)}.$$

But one easily checks (e.g., with Maple) that the maximum of the function on the right hand side in the given region is bounded by 13.622. Hence the statement of the Lemma holds for all $n > 825000 > e^{13.622}$. \diamond

3 Some algorithms for hook lengths

We write H_λ for the multiset of hooklengths of a partition λ , and F_λ for the set of first column hooklengths of λ , and we let h_λ be the product of all the hooklengths of λ .

From now on, λ is always a partition of n , of length l .

The following easy proposition turns out to be very useful.

Proposition 3.1 *Let $p \leq n$ be a prime for which $w_p(\lambda) = \left\lfloor \frac{n}{p} \right\rfloor$. Let μ be the partition obtained from λ by removing its first column. Set $A_\lambda = \{1, \dots, n\} \setminus F_\lambda$. Let A_λ^p and H_μ^p denote the (multi)sets of elements divisible by p in the corresponding sets. Then their (multiset) cardinalities are equal:*

$$|A_\lambda^p| = |H_\mu^p|.$$

Proof. Let F_λ^p denote the set of first column hooklengths of λ divisible by p . By Lemma 2.1 we have

$$\begin{aligned} |F_\lambda^p| + |H_\mu^p| &= w_p(\lambda) = \left\lfloor \frac{n}{p} \right\rfloor \\ &= |\{j \in \{1, \dots, n\} \mid p|j\}| \\ &= |F_\lambda^p \cup A_\lambda^p| = |F_\lambda^p| + |A_\lambda^p| \end{aligned}$$

and hence the stated equality follows. \diamond

Remark. Assume that λ has maximal p -weight for all primes p . Since every “missing” fch of length > 1 has at least one prime factor, the sum of the number of prime factors in all hooklengths of μ should be at least the number of missing fch’s. Indeed, $|A_\lambda|$ is the number of missing fch’s and by the above proposition

$$|A_\lambda| \leq \sum_p |A_\lambda^p| = \sum_p |H_\mu^p|.$$

This is particularly useful in the cases where μ is small.

Corollary 3.2 *Assume that $w_p(\lambda) = \left\lfloor \frac{n}{p} \right\rfloor$ for the prime $p \leq n$.*

- (1) *Let μ be obtained from λ by removing the first column. If $p \nmid h_\mu$, then $p, 2p, \dots, \left\lfloor \frac{n}{p} \right\rfloor p$ are all fch of λ .*
- (2) *If $n - l(\lambda) < p \leq n$, then $p, 2p, \dots, \left\lfloor \frac{n}{p} \right\rfloor p$ are all fch of λ .*
- (3) *Let $1 < j \leq l$. If $h \leq n$ has a prime divisor p satisfying $n - h_j < p$, then h is a fch of λ .*

Proof. Parts (1) and (2) follow immediately from Proposition 3.1.

(3): By Corollary 2.4, all hooks of length divisible by p are in the first column. With μ as in (1), $p \nmid h_\mu$ by Lemma 2.3(2). Hence any multiple of p less than or equal to n is a fch, by (1). In particular, h is a fch of λ . \diamond

Assume now that λ is partition of $n > 4$, of maximal p -weight for all primes $p > 2$.

For all n , we can choose primes p_1, p_2 such that $\frac{n}{2} \leq p_2 < p_1 \leq n$ [3]. By assumption, λ must have hooks of length p_1, p_2 , respectively.

Now by Lemma 2.3 these two hooks are either both in the first row or both in the first column of λ ; w.l.o.g. we assume that they are in the first column,

i.e., they are **fch** (if necessary, we replace λ by its conjugate partition). Then, again by Lemma 2.3, any prime q with $\frac{n}{2} < q \leq n$ is an **fch** of λ . Thus we can deduce the analogue of [1, Proposition 2.12] for our partition λ , i.e.:

Proposition 3.3 *Suppose we have sequences of integers $s_1 < s_2 < \dots < s_r \leq n$, $t_1 < t_2 < \dots < t_r \leq n$ satisfying*

- (i) $s_i < t_i$ for all i ;
- (ii) s_1 and t_1 are primes $> \frac{n}{2}$;
- (iii) For $1 \leq i \leq r - 1$, s_{i+1} and t_{i+1} contain prime factors exceeding $2n - s_i - t_i$.

*Then $s_1, \dots, s_r, t_1, \dots, t_r$ are all **fch** of λ .*

For $n > 3.06 \cdot 10^8$, we know that the algorithm ends close to n , namely, $n - h_1 \leq 225$ [1, Theorem 3.1] (in fact, if n is *sufficiently large*, it arrives at $n - h_1 \leq 2$). Since 225 is too large for our purposes, we have to reduce the bound for $n - h_1$ further by other means. Alternatively, one could also try to prove the tight bound $n - h_1 \leq 3$ for the following improved algorithm.

Proposition 3.4 *Suppose λ is a partition of $n \geq 5$, not a hook, which is of maximal p -weight for all primes $p > 2$. Let $s_1 < s_2 < \dots < s_r \leq n$ be a sequence of integers satisfying*

- (i) $s_1 < s_2$ are **fch** for λ ;
- (ii) for $3 \leq i \leq r$, s_i has a prime divisor exceeding $n - s_{i-2}$.

*Then s_1, \dots, s_r are **fch** of λ .*

Proof. We use induction on i to show that s_i is a **fch**. Let $i \geq 3$ and assume that s_{i-2}, s_{i-1} are **fch**. Since $s_{i-1} > s_{i-2}$ we have $s_{i-2} = h_j$ for some $j > 1$. Then apply Corollary 3.2 with $h = s_i$ to deduce that s_i is a **fch**. \diamond

Note that in the algorithm of Proposition 3.4 we never encounter the case where the prime divisor is 2. It is clearly a strengthening of the algorithm of Proposition 3.3, since it may be applied to each of the sequences s_i and t_i . In particular, any bound for the previous algorithm is also one for this new algorithm.

Example. In the algorithms we typically start with two “large” primes, which are always **fch** of our special partition λ (perhaps after conjugating λ). Take $n = 16$. Choose $s_1 = 11$, $s_2 = 13$. (Note that we have hooks in λ of this length, and – possibly after after conjugation – they are both **fch**.) Our previous algorithm (Proposition 3.3) comes to a halt here. In the new algorithm of Proposition 3.4 we can continue as $s_3 = 14$ has a prime factor $q = 7 > 5 = n - s_1$. Then $s_4 = 15$ has a prime factor $q = 5 > 3 = n - s_2$. As 16 only has the prime factor $q = 2 = n - s_3$, the algorithm terminates here. So the partitions of 16 of maximal weight for all primes > 2 have (up to conjugation) **fch**’s of length 15,14,13,11; it is easy to check that this only holds for the partition (1^{16}) (resp. its conjugate).

Remark 3.5 The algorithm of Proposition 3.4 is very efficient and may quickly be carried out by hand to check that $n - h_1 \leq 3$ holds for all $n \leq 100$. It was also tested with Maple up to $n = 7.5 \cdot 10^8$; it always gave the bound $n - h_1 \leq 3$. Indeed, it almost always ends at $n - h_1 \leq 1$, except for few exceptional values where it ends on the bound 2 and four cases where it ends on 3, namely for 10, 50, 100 and 15.856.204.

4 Proofs of the main results

We first give the proof of Theorem 1.1.

Let $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_l)$ be a partition of n , of length $l = l(\lambda)$. The following *notation* is fixed for λ :

m_1 is the multiplicity of 1 as a part of λ , $k = \lambda_1 - \lambda_2$.

h_1, h_2, \dots, h_l are the **fch** of λ , A_λ is the set of its “missing” **fch**, $d = n - h_1$.

Also, as in the previous section, μ is the partition obtained from λ by removing its first column, i.e. $\mu = (\lambda_1 - 1, \lambda_2 - 1, \dots)$.

Let us assume

(A0) $\lambda \neq (n), (1^n)$.

After possibly replacing λ by its conjugate partition (which has the same p -weight as λ for all p) we are also going to assume

(A1) $l \geq \lambda_1$.

Lemma 4.1 *We have*

(1) $k = \mu_1 - \mu_2 = h_1 - h_2 - 1$.

- (2) $d = |\mu| - \mu_1$ (the depth of μ).
- (3) $h_1(\mu) \leq d + k + 1$.
- (4) $|\mu| \leq 2d + k$.
- (5) $\lambda_1 \leq \frac{h_1+1}{2} \leq l$.
- (6) $h_2 \geq m_1$.
- (7) If $h_2 = m_1$, then λ is a hook partition. Otherwise $h_2 \geq m_1 + 2$.
- (8) $h_2 \geq k$.
- (9) If $h_2 = k$, then $\lambda_1 = l$ and $\lambda_2 = 1$, and hence λ is a hook partition.

Proof. (1) and (2) are trivial. Part (3) follows from the fact that exactly $k + 1$ rim nodes of the (1,1)-hook of μ are in the first row and at most d rim nodes are outside the first row.

Since $\mu_2 \leq d$ we get $\mu_1 \leq d + k$ and thus (4) follows from (3).

Part (5) uses the assumption **(A1)** and the definition of h_1 .

We have $h_2 \geq m_1$ since $\lambda \neq (1^n)$ by **(A0)**; thus (6) holds.

Part (7) is trivial.

For (8) and (9) note that $h_2 = \lambda_2 + l - 2 = \lambda_2 + \lambda_1 + (l - \lambda_1) - 2$, hence $h_2 \geq k$ is equivalent to $(l - \lambda_1) + 2\lambda_2 \geq 2$. Thus the assertion follows easily from **(A1)** and the fact that $\lambda_2 > 0$. \diamond

In addition to **(A0)**, **(A1)**, we now make the assumption

(A2) λ is of maximal p -weight for all primes $p > 2$.

We consider the products

$$\pi_1 := (h_1 + 1)(h_1 + 2) \cdots n, \quad \pi_2 := (h_2 + 1)(h_1 + 2) \cdots (h_1 - 1)$$

having d and k factors, respectively (Lemma 4.1(1)); note that in the case where λ is a hook, $d = 0$, so $\pi_1 = 1$, and thus in this case we will only consider π_2 . By definition, the factors $h_1 + 1, \dots, n$ and $h_2 + 1, \dots, h_1 - 1$ of π_1, π_2 are in A_λ . Thus Proposition 3.1 implies

Lemma 4.2 *If p is an odd prime divisor of π_1 or π_2 , then μ has a hook of length p .*

Let us now deal with the ‘‘hook case’’, i.e., $d = 0$. Because of **(A1)**, the leg of the hook is at least as long as its arm.

Proposition 4.3 *Suppose that $\lambda = (k+1, 1^{n-k-1})$ with $1 \leq k \leq \frac{n-1}{2}$. Then $n = 2^a + 1$ for some $a \in \mathbb{N}$ and $k = 1$, i.e., $\lambda = (2, 1^{n-2})$.*

Proof. By Schur [7] we know that π_2 has a prime divisor $q > k = n - l$. If λ has maximal q -weight, then by Corollary 3.2(2) $\left\lfloor \frac{n}{q} \right\rfloor q$ is a fch of λ . But since $n = h_1 > \left\lfloor \frac{n}{q} \right\rfloor q > h_2$, this is a contradiction. So the only critical case is when $q = 2$ and the 2-weight is non-maximal. But when π_2 has only the prime divisor $q = 2$, then we must have $k = 1$ and $\pi_2 = n - 1$ is a 2-power, as was to be shown. \diamond

Hence we may from now on assume

(A3) λ is not a hook partition, i.e., $d > 0$ and $\lambda_2 > 1$.

We have some further general relations for the parameters of λ :

Lemma 4.4 *We have*

- (1) $k \leq 2d + 1$.
- (2) If $h_1 \geq \frac{n}{2}$ then $d \leq 2k + 1$.
- (3) $|\mu| \leq 4d + 1$.
- (4) If $h_1 \geq \frac{n}{2}$ then $|\mu| \leq 5k + 2$.

Proof. (1) is trivial if $k \leq 1$. If $k > 1$, we may by Lemma 4.1(8) apply Theorem 2.5. It shows that one of the k factors in π_2 has a prime divisor $q > \frac{3}{2}k$, or we have one of the exceptional cases. But none of these can occur:

- (i) $h_2 = 2, h_1 = 5, k = 2$. This is not possible, since by Lemma 4.1(5) $l \geq 3$ and then $h_2 = 2$ implies $\lambda_2 = 1$, a contradiction to **(A3)**.
- (ii) $h_2 = 7, h_1 = 10, k = 2$. We get that (μ_1, μ_2) is (3,1) or (4,2). Both of these partitions are 3-cores, so we get a contradiction to Lemma 4.2.
- (iii) $h_2 = 5, h_1 = 11, k = 5$. Then Lemma 4.1(9) implies that λ has to be a hook, a contradiction to **(A3)**.

As q is now a prime divisor of a number in A_λ , there has to be a hook in μ of length q . (Lemma 4.2.) Thus $q \leq h_1(\mu)$. Using Lemma 4.1(3) we obtain the inequality $\frac{3}{2}k < q \leq h_1(\mu) \leq d + k + 1$, implying $\frac{1}{2}k < d + 1$. Thus certainly $k \leq 2d + 1$. Part (2) is proved in analogy with (1) by applying Theorem 2.5 to the d factors of π_1 . Note that by assumption $d \leq h_1$. Then (3) and (4) follow from (1) and (2), using Lemma 4.1(4). \diamond

We use these relations together with results from [1] and the previous section to reduce $d = n - h_1$ for partitions λ satisfying our assumptions:

Proposition 4.5 *We have $d \leq 4$.*

Proof. When $n \leq 5 \cdot 10^8$, it was already remarked before that with the new algorithm in the previous section we even get down to $d \leq 3$. (We might also use [1] where it was checked that in this range the old algorithm gets down to a bound $d \leq 4$.)

Hence we may now assume that $n > 5 \cdot 10^8$.

Then we obtain $d \leq 225$ using [1, Theorem 3.1] (note that any bound for the old algorithm is also a bound for the new one).

We assume that $5 \leq d \leq 225$, and we want to arrive at a contradiction. By Lemma 4.4(1) we have $k \leq 451$; Lemma 4.4 also implies that the defining factors of π_1 and π_2 are greater than 10^6 . Indeed, these factors are bounded below by $h_2 = n - (d + k + 1) \geq n - (225 + 451 + 1) = n - 677$. We may thus apply Lemma 2.6 and get that π_1 has a prime divisor $q > 2d$. Thus μ must have a hook of length q , by Lemma 4.2. This hook has to be in the first row as there are only $d < q$ nodes in μ below the first row. By Lemma 4.1(3) we have

$$2d < q \leq h_1(\mu) \leq d + k + 1,$$

and thus $d \leq k$.

We may also apply Lemma 2.6 to π_2 and get a prime divisor $q' > 2k \geq 2d$ in one of the k factors in π_2 . Hence μ must have a hook of length divisible by q' in its first row. Moreover, if $q = q'$, then μ must have two q -divisible hook lengths in its first row, by Proposition 3.1. We get

$$2k < q' \leq h_1(\mu) \leq d + 1 + k$$

and thus $k \leq d$. Hence $d = k$. As both q and q' are at least $2d + 1$, one of the two hook lengths is at least $2d + 2$. Thus we obtain

$$2d + 2 \leq h_1(\mu) \leq d + 1 + k,$$

implying $d + 1 \leq k$, a contradiction. \diamond

To finish the proof of Theorem 1.1, we now have only to consider the cases where $d = 1..4$. Here we will see the exceptional cases coming up. First we deal with the case $d = 1$:

Proposition 4.6 *Suppose that $\lambda = (k + 2, 2, 1^{n-k-4})$ with $k \geq 0$. Then $n = 4$ and $\lambda = (2^2)$, $n = 6$ and $\lambda = (3, 2, 1)$ or $n = 8$ and $\lambda = (3, 2, 1^3)$.*

Proof. By Lemma 4.4(1), $k \leq 3$, and by (A1), $n - k - 2 \geq k + 2$, hence $n \geq 2k + 4$. Here, $\mu = (k + 1, 1)$ and $\pi_1 = n$.

If $k = 0$, then $\pi_2 = 1$ and we use the next missing **fch**, i.e., $\pi_3 = n - k - 3 = n - 3$. As $h_\mu = 2$, both n and $n - 3$ have to be 2-powers, hence $n = 4$, and thus $\lambda = (2^2)$. If $k = 1$, then $h_\mu = 3$, and hence $\pi_1\pi_2\pi_3 = n(n - 2)(n - 4) = 2^a3^b$ for some $a, b \in \mathbb{N}$. This is only possible for $n = 6$ and $n = 8$, and in these cases we have $\lambda = (3, 2, 1)$ and $\lambda = (3, 2, 1^3)$, respectively. If $k = 2$, then $h_\mu = 8$, and so $\pi_1\pi_2\pi_3 = n(n - 2)(n - 3)(n - 5)$ has to be a 2-power, which is impossible. If $k = 3$, then $h_\mu = 2 \cdot 3 \cdot 5$; as $\pi_3 = (n - 2)(n - 3)(n - 4)$ is divisible by 3 and has a prime divisor $q \geq 5$ by Theorem 2.5, then both $\pi_1 = n$ and $\pi_2 = n - 6 \geq 4$ have to be 2-powers, which is impossible. \diamond

Next we deal with the case $d = 2$:

Proposition 4.7 *Suppose that $\lambda = (k+3, 3, 1^{n-k-6})$ or $\lambda = (k+2, 2^2, 1^{n-k-6})$ with $k \geq 0$. Then $n = 9$ and $\lambda = (2^3, 1^3)$.*

Proof. Lemma 4.4(1) shows that $k \leq 5$. We have $\mu = (k + 2, 2)$ or $\mu = (k + 1, 1^2)$, respectively. We have $A_\lambda = \{n, n - 1; n - 3, \dots, n - k - 2; n - k - 4, n - k - 5\}$ and $A_\lambda = \{n, n - 1; n - 3, \dots, n - k - 2; n - k - 5\}$, respectively. In particular, the difference between the largest and smallest element is $k + 5 \leq 10$. One can easily check that at most two elements in A_λ are powers of 2 and thus we have at least $k + 2$ (respectively $k + 1$) elements of A_λ divisible by odd primes. Thus μ must have at least $k + 2$ (respectively $k + 1$) hook lengths divisible by an odd prime. The former is not possible at all and the latter only for $k = 0$. Here $h_\mu = 6$ and thus $n(n - 1)(n - 5) = 2^a3^b$ for some $a, b \in \mathbb{N}$. This yields $n = 9$, and then $\lambda = (2^3, 1^3)$. \diamond

Now we discuss the case $d = 3$:

Proposition 4.8 *Suppose λ is one of $(k+4, 4, 1^{n-k-8})$, $(k+3, 3, 2, 1^{n-k-8})$ or $(k+2, 2^3, 1^{n-k-8})$ with $k \geq 0$. Then $n = 10$ and $\lambda = (4, 3, 2, 1)$.*

Proof. Lemma 4.4(1) shows that $k \leq 7$. We have $\mu = (k + 3, 3)$, $\mu = (k + 2, 2, 1)$ or $\mu = (k + 1, 1^3)$, respectively. Moreover, the difference between the largest missing **fch** n and the smallest one $n - k - 7$ is $e = k + 7 \leq 14$. At most three elements in A_λ are powers of 2, since otherwise $n = 16, k = 7$, a contradiction to **(A1)**. Thus we have at least $|\mu| - 3$ odd prime factors in the elements of A_λ . As the hook lengths in μ are at most 11, no hook length has two different odd prime divisors. Thus μ must have at least $|\mu| - 3$ hook lengths divisible by an odd prime. This eliminates all possibilities for μ except $(3, 2, 1)$, $(2, 1^3)$ and (1^4) ; note that in all of these cases the elements

in A_λ are not divisible by primes > 5 . As $\pi_1 = n(n-1)(n-2)$ has a prime divisor $q \geq 5$ by Theorem 2.5, $\mu = (1^4)$ cannot occur. For $\mu = (2, 1^3)$ we have $h_\mu = 2 \cdot 3 \cdot 5$, hence as π_1 is divisible by 3 and 5, the other elements $n-4$ and $n-8$ in A_λ have to be 2-powers; but this is impossible. If $\mu = (3, 2, 1)$, then $A_\lambda = \{n, n-1, n-2, n-4, n-6, n-8\}$. A parity consideration shows that this is only possible for $n = 10$. \diamond

Proposition 4.9 *The case $d = 4$ is not possible.*

Proof. By Remark 3.5 we may assume that n is large so that we may apply Lemma 2.6 to π_2 , if $k \neq 0, 1, 3, 4$. We conclude as before that μ must have a hook divisible by a prime $q > 2k$ so that $2k < q \leq h_1(\mu) \leq d + k + 1$ yielding $k \leq d = 4$. We thus have in any case $k \leq 4$ and hence $|\mu| \leq 12$ by Lemma 4.1. Applying Theorem 2.5 to π_1 we see that μ must have a hook of length 7 or 11. If also $k = 4$ then Theorem 2.5 applied to π_2 shows that μ has another hook of length 7 or 11. But these both have to be in the first row of μ , and this is easily seen not to be possible. If $k = 2, 3$ then Theorem 2.5 shows that μ has another hook of length divisible by 5. This is only possible for $\mu = (7, 4)$, $\mu = (6, 4)$ or $\mu = (5, 3, 1)$. If $k \leq 1$ then $h_1(\mu) \leq 6$, a contradiction. As π_1 is divisible by 3, but $\mu = (5, 3, 1)$ is a 3-core, this is excluded. For the other two possibilities, apply Theorem 2.5 to $\pi_3 = (n-k-6)(n-k-7)(n-k-8)(n-k-9)$ to see that μ must have a further hook of length divisible by 7 or 11, which is not possible. \diamond

Now the proof of Theorem 1.1 is complete, and with this result at hand, we may now prove Theorem 1.2 and Theorem 1.5.

Proof of Theorem 1.2. First, take a non-linear irreducible character χ of S_n ; thus, $\chi = [\lambda]$, where λ is a partition of n different from $(n), (1^n)$. As $[2^2]$ is zero on transpositions, and $[2^2](1) = 2$, we may now assume that $\lambda \neq (2^2)$. Hence by Theorem 1.1(1) there exists a prime p , such that $w_p(\lambda) < w = \left\lfloor \frac{n}{p} \right\rfloor$. By the Murnaghan-Nakayama formula ([4, 2.4.7], 2.4.7), the character then vanishes on a product of w p -cycles. Indeed, by Theorem 1.1(2) we may choose $p > 2$ except for the partitions occurring in this case; we deal with these below to check the final assertion.

Now consider a non-linear irreducible character of A_n , labelled by λ , hence $\lambda \neq (n), (1^n), (2, 1), (2^2)$. If λ is not of maximal p -weight for some $p > 2$, then χ vanishes on the same element σ of order p as $[\lambda]$ (note that $\sigma \in A_n$).

Thus, the only critical characters are the ones labelled by the exceptional partitions listed in Theorem 1.1(2), which are of non-maximal p -weight only for $p = 2$. Of course, it suffices to consider one of a pair of conjugate partitions.

Now, if $n = 2^a + 1$, $a > 1$, then $(n - 1, 1)$ is of 2-weight $2^{a-1} - 1$, and the character $[n - 1, 1]$ and its irreducible restriction to A_n vanish on a product of 2^{a-1} transpositions (which is an element of A_n); note that $[n - 1, 1]$ is of 2-power degree 2^a . The irreducible characters of A_6 (and S_6) labelled by $(3, 2, 1)$ are zero on elements of cycle type $(2^2, 1^2)$ (these characters are of degree 8 and 16, respectively). The character $[5, 2, 1]$ and its irreducible restriction to A_8 is zero on elements of cycle type $(2^2, 1^4)$ (its degree is 64), and $[6, 3]$ and its irreducible restriction to A_9 is zero on elements of cycle types $(2^4, 1)$ and $(3^2, 1^3)$. Finally, the irreducible characters of A_{10} (and S_{10}) labelled by $(4, 3, 2, 1)$ are zero on elements of cycle types $(2^4, 1^2)$, $(2^2, 1^6)$ and $(3^2, 1^4)$. \diamond

Proof of Theorem 1.5. (1). The first assertion is immediate from Theorem 1.1. The second follows from the first, using that for $n \notin \{1, 2, 3, 4, 6\}$ there exist primes p such that $\frac{n}{2} < p < n - 1$ to see that (1^n) is not in the principal p -block.

(2) This follows similarly as (1). \diamond

Acknowledgement. The authors would like to thank the Isaac Newton Institute for Mathematical Sciences of the University of Cambridge for its hospitality during a stay there in the frame of the programme *Symmetric functions and Macdonald polynomials*, where some part of the work for this article was done.

References

- [1] A. Balog, C. Bessenrodt, J. B. Olsson, K. Ono, Prime power degree representations of the symmetric and alternating groups, *J. London Math. Soc.* (2) 64 (2001) 344-356
- [2] D. Hanson, On a theorem of Sylvester and Schur, *Canad. Math. Bull.* 16 (1973) 195-199
- [3] H. Harborth, A. Kemnitz, Calculations for Bertrand's postulate, *Math. Mag.* 54 (1981) 33-34

- [4] G. James, A. Kerber, *The representation theory of the symmetric group*. Encyclopedia of Mathematics and its Applications, 16, Addison-Wesley 1981
- [5] G. Malle, G. Navarro, J.B. Olsson, Zeros of characters of finite groups, *J. Group Theory* 3 (2000) 353-368
- [6] J.B. Olsson, *Combinatorics and representations of finite groups*, Vorlesungen aus dem FB Mathematik der Univ. Essen, Heft 20, 1993
- [7] I. Schur, Einige Sätze über Primzahlen mit Anwendungen auf Irreduzibilitätsfragen. I. *Sitzungsberichte der Preuss. Akad. d. Wiss.* 1929, Phys.-Math. Klasse, 125-136. (Collected works, vol 3, 140-151)

Christine Bessenrodt

*Fachbereich Mathematik, Universität Hannover
Welfengarten 1, D-30167 Hannover, Germany
Email address: bessen@math.uni-hannover.de*

Jørn B. Olsson

*Matematisk Institut, Københavns Universitet
Universitetsparken 5, DK-2100 Copenhagen Ø, Denmark
Email address: olsson@math.ku.dk*