

Basis Expansions

With $X \in \mathbb{R}^p$ and $Y \in \mathbb{R}$ the function

$$f(x) = E(Y|X = x)$$

is typically globally a non-linear function. We discuss situations where p is small or moderate, but where the function is complicated.

A **basis function expansion** of f is an expansion

$$f(x) = \sum_{m=1}^M \beta_m h_m(x)$$

with $h_m : \mathbb{R}^p \rightarrow \mathbb{R}$ for $m = 1, \dots, M$.

The basis functions are chosen and fixed and the parameters β_m for $m = 1, \dots, M$ are estimated. This is a **linear model** in the **derived variables** $h_1(X), \dots, h_M(X)$.

Polynomial Bases

Monomials are classical basis functions;

$$h_m(x) = x_1^{r_1} x_2^{r_2} \dots x_p^{r_p}$$

with $r_i \in \{0, \dots, d\}$ and $r_1 + \dots + r_p \leq d$. This basis spans the polynomials of degree $\leq d$.

- If the linear models provide first order Taylor approximations of the function, expansions in the degree d polynomials provide order d Taylor approximations.
- However, if $p \geq 2$ the number of basis functions grows exponentially in d .

Indicators

A completely different, non-differentiable idea is to approximate f locally as a constant. Box-type basis functions are

$$h_m(x) = 1(l_1 \leq x_1 \leq r_1) \dots 1(l_p \leq x_p \leq r_p)$$

with $l_i \leq r_i$ and $l_i, r_i \in [-\infty, \infty]$ for $i = 1, \dots, p$.

If the boxes are disjoint, the columns in the \mathbf{X} -matrix for the derived variables are orthogonal:

$$\mathbf{X}_{im} = h_m(x_i) \in \{0, 1\}$$

We can think of this as **dummy variables representing the box**. Consequently, with least squares estimation

$$\hat{\beta}_m = \frac{1}{N_m} \sum_{i: h_m(x_i)=1} y_i, \quad N_m = \sum_{i=1}^N 1(h_m(x_i) = 1).$$

Basis Strategies

The size of the typical set of basis functions increases rapidly with p .
What are feasible strategies for basis selection?

- **Restriction:** Choose a priori only special basis functions
 - Additivity; $h_{mj} : \mathbb{R} \rightarrow \mathbb{R}$

$$h_m(x) = \sum_{j=1}^p h_{mj}(x_j)$$

- Radial basis functions:

$$h_m(x) = D \left(\frac{\|x - \xi_m\|}{\lambda_m} \right)$$

- **Selection:** As variable selection – implement exhaustive or step-wise inclusions/exclusions of basis functions.
- **Penalization:** As ridge regression – keep the entire set of basis functions but penalize the size of the parameter vector.

Figure 5.1

Figure 5.2

Splines – $p = 1$

Define $h_1(x) = 1$, $h_2(x) = x$ and

$$h_{m+2}(x) = (x - \xi_m)_+ \quad t_+ = \max\{0, t\}$$

for ξ_1, \dots, ξ_K the **knots**.

$$f(x) = \sum_{m=1}^{M+2} \beta_m h_m(x)$$

is a **piecewise linear, continuous** function. One **order- M spline basis** with knots ξ_1, \dots, ξ_K is

$$h_1(x) = 1, \dots, h_M(x) = x^{M-1}, \quad h_{M+l}(x) = (x - \xi_l)_+^{M-1}, \quad l = 1, \dots, K.$$

Figure 5.3

Natural Cubic Splines

Splines of order M are polynomials of degree $M - 1$ beyond the boundary knots ξ_1 and ξ_K . The **natural cubic splines** are the splines of order 4 that are linear beyond the two boundary knots. With

$$f(x) = \beta_0 + \beta_1 x + \beta_2 x^2 + \beta_3 x^3 + \sum_{k=1}^K \theta_k (x - \xi_k)_+^3$$

the restriction is that $\beta_2 = \beta_3 = 0$ and

$$\sum_{k=1}^K \theta_k = \sum_{k=1}^K \theta_k \xi_k = 0.$$

$$N_1(x) = 1, \quad N_2(x) = x$$

and

$$N_{2+l}(x) = \frac{(x - \xi_l)_+^3 - (x - \xi_K)_+^3}{\xi_K - \xi_l} - \frac{(x - \xi_{K-1})_+^3 - (x - \xi_K)_+^3}{\xi_K - \xi_{K-1}}$$

for $l = 1, \dots, K - 2$ form a basis.

B Splines (Basis Splines, or YASB)

Yet Another Spline Basis ...

Defined by a recursion in M ;

$$B_{k,1}(x) = \begin{cases} 1 & \text{if } \tau_k \leq x \leq \tau_{k+1} \\ 0 & \text{otherwise} \end{cases}$$

with

$$\tau_1 \leq \dots \tau_M = \xi_0 < \tau_{M+1} = \xi_1 < \dots < \tau_{M+K} = \xi_K < \tau_{M+K+1} = \xi_{K+1} \leq \dots \leq \tau_{2M+K}$$

and

$$B_{k,r} = \frac{x - \tau_i}{\tau_{i+r+1} - \tau_i} B_{k,r-1}(x) + \frac{\tau_{i+r} - x}{\tau_{i+r} - \tau_i} B_{k+1,r-1}(x)$$

for $k = 1, \dots, K + 2M - r$.

Figure 5.20 – B-splines

Knot Placing Strategies

How do you determine the knots?

- Fix the number (the complexity parameter), spread them uniformly over the whole range of data.
- Fix the number, spread them according to the empirical distribution.
- Adaptive selection of the number and/or the location – ranging from ad hoc adaptation to a full fledged, complete estimation from data.
- Smoothing methods automatically determine their location

Smoothing Splines

Allowing $E(Y|X = x) = f(x)$ to be an arbitrary, but twice differentiable function, define the **penalized residual sum of squares**

$$\text{RSS}(f, \lambda) = \sum_{i=1}^N (y_i - f(x_i))^2 + \lambda \int_a^b f''(t)^2 dt$$

If f^λ is a minimizer of $\text{RSS}(f, \lambda)$, the **natural cubic splines** with knots in x_1, \dots, x_N have the properties that

- they can interpolate; there is a natural cubic spline f_0 with $f_0(x_i) = f^\lambda(x_i)$
- and among all interpolants f , f_0 attains the smallest value of

$$\int_a^b f''(t)^2 dt.$$

The solution $f^\lambda = \sum_{i=1}^N \theta_i N_i(x)$ is a natural cubic spline.

Smoothing Splines

In vector notation

$$\mathbf{f} = \mathbf{N}\theta$$

with $\mathbf{N}_{ij} = N_j(x_i)$ and

$$\begin{aligned}\text{RSS}(f, \lambda) &= (\mathbf{y} - \mathbf{f})^T (\mathbf{y} - \mathbf{f}) + \lambda \int_a^b f''(t)^2 dt \\ &= (\mathbf{y} - \mathbf{N}\theta)^T (\mathbf{y} - \mathbf{N}\theta) + \lambda \theta^T \mathbf{\Omega}_N \theta\end{aligned}$$

with

$$\mathbf{\Omega}_{N,ij} = \int_a^b N_i''(t) N_j''(t) dt.$$

This **generalized ridge regression problem** has solution

$$\hat{\theta} = (\mathbf{N}^T \mathbf{N} + \lambda \mathbf{\Omega}_N)^{-1} \mathbf{N}^T \mathbf{y}$$

and the fitted values are

$$\hat{\mathbf{f}} = \mathbf{N}(\mathbf{N}^T \mathbf{N} + \lambda \mathbf{\Omega}_N)^{-1} \mathbf{N}^T \mathbf{y}$$

Degrees Of Freedom

Writing

$$\mathbf{S}_\lambda = \mathbf{N}(\mathbf{N}^T \mathbf{N} + \lambda \mathbf{\Omega}_N)^{-1} \mathbf{N}^T$$

and by analogy with projection matrices the **effective degrees of freedom** is

$$\text{df}_\lambda = \text{trace}(\mathbf{S}_\lambda).$$

The value of df_λ is monotonely decreasing from N to 0 as λ increases from 0 to ∞ .

The matrix \mathbf{S}_λ is known as a **spline smoother** and it is common to specify the degrees of freedom instead of λ in practice.

Figure 5.8 – Smoother Matrix

Multidimensional Splines

Two multivariate versions.

- **Tensor products.** Consider a basis consisting of

$$B_{i_1,R}(x_1)B_{i_2,R}(x_2)\dots B_{i_p,R}(x_p)$$

– compare with the multinomial basis for polynomials.

- **Thin plate splines.** If $p = 2$ consider minimizing

$$\sum_{i=1}^N (y_i - f(x_i))^2 + \lambda \int_A (\partial_1^2 f)^2 + 2(\partial_1 \partial_2 f)^2 + (\partial_2^2 f)^2.$$

The solution is a function

$$f(x) = \beta_0 + x^T \beta + \sum_{i=1}^N \alpha_i \eta(\|x - x_i\|)$$

with $\eta(z) = z^2 \log(z^2)$ – thus a **radial basis function expansion**.

Figure 5.10 – Tensor Products of B-splines